



You have a

Maisonneuve Injury

This is an injury to ligaments in the lower leg called the syndesmosis and may also be combined with a fracture.

Healing:

It can take up to eight weeks for the first symptoms to settle.

Smoking will slow down your healing. We would advise that you stop smoking while your fracture heals. Talk to your GP or go to www.smokefree.nhs.uk for more information.

Pain and swelling:

Your ankle may be swollen and painful. Swelling is often worse at the end of the day. It is normal to have mild pain and swelling for 6-12 months after your injury.

Taking pain medication, elevating your ankle and using ice or cold packs will help. More information is on the next page.



Walking and your boot:

When standing and walking you should only lightly touch your foot to the floor. You should not put weight through your foot.

The boot protects your ankle and will make you more comfortable. Wear the boot when you are standing and walking for the first six weeks. You can take it off at night and at rest. Please inform us if you are diabetic; you may need a special boot.

You are allowed to put weight through your foot. You may find it easier to use crutches in the early stages.



Exercises:

It is important to start exercises as soon as possible. Instructions are on the next page.

Follow up:

There is a small chance that this injury can be unstable which may require an operation. You will have a weight bearing x-ray and see a specialist normally approximately two weeks after your injury. The specialist will explain the results and the next stage of your care.

Any questions:

If you are concerned about your symptoms, unable to follow this rehabilitation plan or have not received your appointment letter please contact us.



Caring for your injury: Week 1-6

Wear your boot whenever standing and walking for the first six weeks. Remember to only touch your foot to the floor for balance; you must not put weight through the foot. The specialist will tell you when you can begin to put more weight through your foot.

You can remove the boot when resting and at night. Wear a long sock in your boot.

Using a cold pack will help with your pain and swelling. You can use an ice pack or bag of frozen peas wrapped in a damp towel. Put this on your ankle for up to 15 minutes every few hours. Make sure the ice isn't in direct contact with your skin.

Try to rest your ankle, especially in the first 24-72 hours. Raise your ankle on a stool or cushions so that it is above the level of your hip. This will help to reduce your swelling.


Exercises

Early movement of the ankle and foot is important to promote circulation and reduce the risk of developing a Deep Vein Thrombosis (blood clot).

Do these exercises three to four times a day. Start straight away, working within your pain levels.



1. Point your foot up and down. Repeat this 10 times.



2. With your heels together, move your toes apart to turn the foot outwards. Repeat this 10 times. Do this movement gently within comfort.



3. Make gentle circles with your foot in one direction and then the other direction. Repeat this 10 times.



Caring for your injury: Week 6-12

You can now stop using your boot. Start by walking without the boot around your own home. Build up to walking without it outside or for longer walks. You should no longer need the boot or crutches after 8 weeks post injury.

It is normal to still have mild discomfort and swelling. This may continue for six to twelve months.

Activity and Exercise

Gradually increase your level of activity. You should avoid impact activity for three months. This includes running, jumping and dancing.

You can now progress your exercises.

Stretches



1. Sit with your leg straight out in front of you. Put a towel or bandage around your foot and pull it towards you. Feel a stretch in the back of the calf. Hold for 30 seconds.




2. Point your toes down as far as they go, place your other foot on top and apply some pressure. This will stretch the top of your foot. Hold for 30 seconds.



Balance

Level 1: These exercises are for people who couldn't stand on one leg before their injury.

1. Stand with your feet as close together as possible. Hold onto a firm support in safe space. Hold your balance for 30 seconds.
2. Now try removing your hand. Try to keep your balance for 30 seconds.
3. Hold onto a firm support. Put one foot in front of the other, as close together as you can. Hold this for 30 seconds. If you can, try to let go of the support and keep your balance.



Level 2: These exercises are for people who could stand on one leg before their injury.

1. Hold onto a firm surface in a safe space. Try to stand on one leg. Hold this for 30 seconds. Stop if you experience pain. When you can do this comfortably, try the next exercise.
2. Try to stand on one leg without holding on to a support. Try to hold this for 30 seconds. When you can do this comfortably, try the next exercise.
3. You can try these exercises with your eyes closed. Make sure you are always in a safe environment with a support to hold if needed.



Contact the Virtual Fracture Clinic if you are struggling to recover your movement or return to activity.



Frequently Asked Questions

I am struggling with my boot. What do I do?

The boot has a thicker sole; this can make you feel uneven. Make sure you wear a supportive shoe or trainer on your uninjured foot. This will reduce stress on other joints.

If you need more advice contact the Virtual Fracture Clinic.

I am diabetic, does this change things?

If you are diabetic please contact us to discuss your boot. This is particularly important if you have problems with your skin or sensation. We may provide you with a specialist diabetic boot.

When can I start driving?

You can return to driving when:

- You are no longer using your boot,
- You can walk comfortably and
- You can perform an emergency stop pain free.

Always test your ability to drive in a safe environment first.

How can I get a certificate for work?

You can get a fitness for work statement from your GP or the doctor at your Fracture Clinic appointment.

What do I do with my boot and crutches when I no longer need them?

We are not able to use boots again. These should not be return to the hospital.

Crutches can be returned to the Fracture Clinic or A&E.

How do I contact the Virtual Fracture Clinic?

Call 020 7288 3310



Patient advice and liaison service (PALS)

If you have a compliment, complaint or concern please contact our PALS team on 020 7288 5551 or whh-tr.whitthealthPALS@nhs.net

If you need a large print, audio or translated copy of this leaflet please email whh-tr.patient-information@nhs.net. We will try our best to meet your needs.

[Twitter.com/WhitHealth](https://twitter.com/WhitHealth)
[Facebook.com/WhittingtonHealth](https://facebook.com/WhittingtonHealth)

Whittington Health NHS Trust
Magdala Avenue
London
N19 5NF
Phone: 020 7272 3070
www.whittington.nhs.uk

Date published: 07/02/2022
Review date: 07/02/2024
Ref: SC/T&O-VFC/MNI/01

© Whittington Health
Please recycle

