



You have an

Midfoot Fracture – NWB

This is a break to a bone in the middle of your foot.

Healing:

It can take up to 6-12 weeks for this fracture to heal.

Smoking will slow down your healing. We would advise that you stop smoking while your fracture heals. Talk to your GP or go to www.smokefree.nhs.uk for more information.

Pain and swelling:

You may have foot pain and swelling for 3-6 months after your injury. Swelling is often worse at the end of the day.

Taking pain medication, elevating your foot and using ice or cold packs will help. More information is on the next page.



Walking and your boot:

You are only allowed to touch your foot on the floor for balance. You should not put weight through your foot; it should rest lightly on the floor only. You will need crutches to help you walk.

The boot protects your foot and will make you more comfortable. Wear the boot when you are standing and walking. You can take it off at night and at rest. You need to wear the boot for at least six weeks after your injury. Please inform us if you are diabetic; you may require a specialist boot.



Exercises:

It is important to start exercises as soon as possible. Instructions are on the next page.

Follow up:

A follow up appointment is not normally needed for this injury. If you still have significant pain and swelling after three months, then please contact the Virtual Fracture Clinic team.

Any questions:

If you are concerned about your symptoms, are unable to follow this rehabilitation plan or have pain other than at the site of your injury please contact the Virtual Fracture Clinic team.



Caring for your injury: Week 1-2

Remember to wear your boot whenever standing and walking. You should only rest your foot on the floor lightly. You will need to walk like this for the first two weeks.

You can remove the boot when resting, at night and to wash and dress. Wear a long sock in your boot.

Using a cold pack will help with your pain and swelling. You can use an ice pack or bag of frozen peas wrapped in a damp towel. Put this on your foot for up to 15 minutes every few hours. Make sure the ice is not in direct contact with your skin.

Try to rest your foot, especially in the first 24-72 hours. Raise your foot on a stool or cushions so that it is above the level of your hip. This will help to reduce your swelling.


Exercises:

Early movement of the ankle and foot is important to promote circulation and reduce the risk of developing a Deep Vein Thrombosis (blood clot).

Do these exercises 3-4 times a day. Start straight away, you do not need to push into pain.



1. Point your foot up and down.
Repeat this 10 times.



2. With your heels together, move your toes apart to turn the foot outwards.
Repeat this 10 times. Do this movement gently within comfort.



3. Make gentle circles with your foot in one direction and then the other direction.
Repeat this 10 times.



Caring for your injury: Week 2-6

It is now two weeks since your injury and you can start to put more weight through your foot. Keep using your boot when you are standing and walking. By six weeks after your injury you should be able to put full weight through your foot. You can stop using your crutches as soon as you feel ready.

It is normal to still have mild discomfort and swelling. This may continue for three to six months. Keep elevating your leg and using ice packs.

Activity and Exercise

Keep doing your exercises until you have full movement in your foot.

Caring for your injury: Week 6-12

It is now 6 weeks since your injury and you can start to wean out of you boot. Stop using it around your own home first; you may still need it for longer walks. You should aim to be walking without your boot by six to eight weeks after injury.

It is normal to still have mild discomfort and swelling. This may continue for three to six months.

Activity and Exercise

Keep doing your exercises until you have full movement in your foot.

Gradually increase your level of activity. You should avoid impact activity for three months. This includes running, jumping and dancing.

Frequently Asked Questions

I am struggling with my boot. What do I do?

The boot has a thicker sole; this can make you feel uneven. Make sure you wear a supportive shoe or trainer on your uninjured foot. This will reduce stress on other joints.

If you need more advice contact the Virtual Fracture Clinic.

I am diabetic, does this change things?

If you are diabetic please contact us to discuss your boot. This is particularly important if you have problems with your skin. We may provide you with a specialist diabetic boot.



When can I start driving?

You can return to driving when:

- You are no longer using your boot,
- You can walk comfortably
- You can perform an emergency stop pain free.

Always test your ability to drive in a safe environment first.

How can I get a certificate for work?

You can get a fitness for work statement from your GP or the doctor at your Fracture Clinic appointment.

What do I do with my boot and crutches when I no longer need them?

We are not able to use boots again. These should not be returned to the hospital.

Crutches can be returned to the Fracture Clinic or A&E.

How do I contact the Virtual Fracture Clinic?

Call 020 7288 3310

With thanks to Brighton and Sussex University Hospitals NHS Trust for kindly sharing their information sheets with us with full permission to use contents and images.

Patient advice and liaison service (PALS)

If you have a compliment, complaint or concern please contact our PALS team on 020 7288 5551 or whh-tr.whitthealthPALS@nhs.net

If you need a large print, audio or translated copy of this leaflet please email whh-tr.patient-information@nhs.net. We will try our best to meet your needs.

[Twitter.com/WhitHealth](https://twitter.com/WhitHealth)

[Facebook.com/WhittingtonHealth](https://facebook.com/WhittingtonHealth)

Whittington Health NHS Trust

Magdala Avenue

London

N19 5NF

Phone: 020 7272 3070

www.whittington.nhs.uk

Date published: 19/04/2022

Review date: 19/04/2024

Ref: SC/T&O-VFC/MFF-NWB/01

© Whittington Health

Please recycle

