



North London Obesity Surgery Service

Vitamin B12

Patient Information

What does vitamin B12 do?

Vitamin B12 is one of the B-complex vitamins. It is important for the function of your nerves and for the production of the DNA and RNA in your cells. It also works together with folic acid to make red blood cells and other compounds that are important for your cardiovascular and immune systems.

What happens if my B12 is low / if I do not get enough B12?

It is common for people with low B12 to develop anaemia because they cannot make blood cells normally. With this anaemia, cells become large and the nucleus of the cell is not formed properly. If B12 deficiency is allowed to continue for a long time, serious problems can develop such as permanent damage to the nerves, memory loss and dementia.

If you do not have enough B12 you may notice that you become fatigued easily and weak. You may experience dizziness, heart palpitations (rapid heartbeats), and feel short of breath.

Other common symptoms are poor appetite, sore/swollen tongue, and numbness and tingling of the handsand feet.

What are other symptoms?

- Diarrhoea
- Yellowish skin and eyes
- Irritability and depression
- Psychosis
- · Muscle weakness, difficulty walking
- Changes in vision
- · Burning sensation around mouth

Where can I get B12?

B12 is only found in foods of animal origins. Plants do not make B12 – so fruits, vegetables, beans and grains will not supply this vitamin. Good dietary sources of vitamin B12 include fish, dairy products, organmeats (particularly liver and kidney), eggs, beef, and pork. You can also get B12 in dietary supplements.

It is found in most multivitamins, in B-complex, and alone in tablets, capsules, liquids and sublinguals (pills that dissolve under the tongue). It is also available as a prescription as injections or as a nasal spray. If you have B12 deficiency, your doctor will probably ask you to take supplemental or prescription B12 and watchyour dietary intake.



Other important things

If you have weight loss surgery such as gastric bypass, it is harder for the body to absorb B12 from food and from some types of supplements. For this reason, your doctor may ask you to use a sublingual product, or to get regular injections to prevent problems. If you have B12 deficiency, it is important that you follow up with your doctor for lab tests and other recommended care.

> Should you need more information, please contact our Specialist Bariatric Dietitians on 0207 288 5989.

Patient advice and liaison service (PALS)

If you have a compliment, complaint or concern please contact our PALS team on 020 7288 5551 or whh-tr.PALS@nhs.net

If you need a large print, audio or translated copy of this leaflet please email whh-tr.patient-information@nhs.net. We will try our best to meet your needs.

Twitter.com/WhitHealth Facebook.com/WhittingtonHealth

Whittington Health NHS Trust Magdala Avenue London N19 5NF Phone: 020 7272 3070

www.whittington.nhs.uk

Date published: 30/08/2022 Review date: 30/08/2024 Ref: SC/Bariatr/VitB12/01

© Whittington Health

Please recycle

